arXiv:cond-mat/9510086v1 16 Oct 1995

Skyrmions and edge spin excitations in quantum Hall droplets

J. H. Oaknin[1], L. Martín-Moreno[2] and C. Tejedor[1]
[1]*Departamento de Física de la Materia Condensada, Universidad Autónoma de Madrid, Cantoblanco, 28049, Madrid, Spain.*
[2]*Instituto de Ciencia de Materiales (CSIC), Universidad Autónoma de Madrid, Cantoblanco, 28049, Madrid, Spain.*

(February 6, 2008)

We present an analysis of spin-textures in Quantum Hall droplets, for filling factors $\nu \simeq 1$. Analytical wavefunctions with well defined quantum numbers are given for the low-lying states of the system which result to be either bulk skyrmions or edge spin excitations. We compute dispersion relations and study how skyrmions become ground states of the Quantum Hall droplet at $\nu \gtrsim 1$. A Hartree-Fock approximation is recovered and discussed for those spin textures.

PACS number: 73.40.Hm

Spin excitations of a two-dimensional (2D) electron gas in the filling factor $\nu = 1$ regime are the subject of increasing interest due to the experimental observation [1–3] of skyrmions. [4–9] They have been analyzed both by a classical nonlinear-σ model (CNLSM) [5,7] and by a Hartree-Fock (HF) approximation. [6,9] These descriptions ignore quantum fluctuations; furthermore quantum numbers as modulus (S^2) and third component (S_z) of the spin, and total (M) and center-of-mass (M_{CM}) angular momenta are not well described. A superposition of mean-field wavefunctions has been proposed [10] which have well defined M and S_z but still neither M_{CM} nor S^2 are represented adequately. Current descriptions of spin-textures have always been considered in infinite [5–7,9] or periodic (sphere) [4,7,10] systems. Therefore, they can not be directly used to analyze available experimental information for small systems. [11–13]

In this letter we address the two previously mentioned restrictions of present theories:

1) We analyze spin-textures in a system of N electrons moving in 2D, in the presence of both a high perpendicular magnetic field B and a confinement potential (Quantum Hall droplets (QHD)).

2) We obtain analytical many-body wavefunctions of spin excitations with all quantum numbers (M, M_{CM}, S^2 and S_z) properly described, i.e. we present a microscopic description including quantum fluctuations.

We also obtain the energies of these excitations, which turn out to be low-lying excitations at $\nu = 1$, some of them becoming the ground state (GS) close to $\nu = 1$.

We consider interacting electrons moving in the xy plane, confined by a parabolic potential (characterised by a bare frequency ω_0). We assume that B is high enough as to project the Hamiltonian H onto the lowest Landau level with both spin up and down. In symmetric gauge,

$$H = H^{SP} + \sum_{m_i,\sigma_j} \frac{V_{m_1 m_2 m_3 m_4}}{2} c^\dagger_{m_1,\sigma_1} c^\dagger_{m_2,\sigma_2} c_{m_3,\sigma_2} c_{m_4,\sigma_1} \qquad (1)$$

where $c^\dagger$ and c are the electron creation and annihilation operators, respectively, m_i are single-particle (SP) angular momenta and σ_j label spins. We consider matrix elements $V_{m_1 m_2 m_3 m_4}$ for the Coulomb interaction, although we have checked that our conclusions do not depend on the details of the electron-electron repulsion. The single particle part H^{SP} has a spectrum

$$E^{SP}(M, S_z) = \left(\frac{N}{2} + \frac{\Omega - \omega_c}{2\Omega} M \right) \hbar\Omega + g\mu_B S_z B \qquad (2)$$

where $\omega_c = eB/(m^* c)$, $\Omega = [\omega_c^2 + 4\omega_0^2]^{1/2}$, g is the Lande factor and μ_B the Bohr magneton. Due to the symmetry of the system the spectrum separates in subspaces with well defined M, M_{CM}, S^2, S_z. The energy ordering of states within a subspace does not depend neither on ω_0 nor on B; only the relative positioning of energies in different subspaces do.

Some of these subspaces have dimension 1, so the state which spans each one of them is an eigenstate of the many-body Hamiltonian. This is the case for single Slater determinants comprised of the k lowest m SP states with spin $\downarrow$ and the $N - k$ lowest m SP states with spin $\uparrow$, where $k = 0, 1, ...N/2$. We label these eigenstates as $|C^k_{N-k}\rangle$, and we refer to them as "compact" states, due to the the fact that they have minimum M compatible with a given $S_z = (N-2k)/2$. Numerical calculations [14] show that compact states are GS for some ranges of the parameters defining the QHD. For example, on increasing B the transition from the paramagnetic $|C^{N/2}_{N/2}\rangle$ ($\nu = 2$) to the ferromagnetic $|C^0_N\rangle$ ($\nu = 1$) occurs by flipping spins through all $|C^k_{N-k}\rangle$. However, these are not the only GS in the process, other GS with small S_z appearing in between any spin-flip transition. [14] Small-S_z states appear as well as GS for $\nu \lesssim 1$. [15,16]

We want to describe these highly correlated low-S_z GS and, more generaly, low-lying excitations close to $\nu = 1$. It has been shown that there is a branch of low-lying excitations at $\nu = 1$ which resemble magnetoexcitons of $|C^0_N\rangle$. [17] By analogy, in order to describe spin-charge excitations, we try the ansatz of magnon-like excitations: $\mathcal{O}^\dagger_{\pm\Delta}|C^k_{N-k}\rangle$, with $\mathcal{O}^\dagger_{\pm\Delta} = \sum_m o_m c^\dagger_{m\pm\Delta,\downarrow} c_{m,\uparrow}$. It is known that low-lying excitations have $M_{CM} = 0$. [18] As for charge excitations at $\nu = 1$, [17] imposing this condition fixes the values of the coefficients o_m. We find [19] several possibilities for $\mathcal{O}^\dagger_{\pm\Delta}$ (with $\Delta > 0$):

$$\Upsilon^\dagger_{-\Delta} = \sum_m \sqrt{\frac{(m+\Delta)!}{m!}} c^\dagger_{m-\Delta,\downarrow} c_{m,\uparrow} \qquad (3)$$

$$\Lambda^{\dagger}_{\Delta} = \sum_{m} \sqrt{\frac{m!}{(m+\Delta)!}}\, c^{\dagger}_{m+\Delta,\downarrow} c_{m,\uparrow} \tag{4}$$

It can be shown that $(\mathcal{O}^{\dagger}_{\pm\Delta})^n$, n being an integer and $\mathcal{O}^{\dagger}_{\pm\Delta}$ any of the previous operators, also generate states with well defined S_z, M, and M_{CM}. For that purpose, these operators can act on *any* state $|C^k_{N-k}\rangle$, except $(\Lambda^{\dagger}_{\Delta})^n$, which requires $\Delta \leq k$ (note that therefore none of the $(\Lambda^{\dagger}_{\Delta})^n$ act on the $\nu = 1$ state). However, $(\mathcal{O}^{\dagger}_{\pm\Delta})^n$ generate states without a well defined S^2. Due to the Zeeman term, the lowest energy state in a (M, S^2) subspace has maximum possible S_z. Let us call $\mathcal{P}$ the operator which projects onto the subspace $S^2 = S_z(S_z + 1)$. [20] So we generate, from compact states, sets of states $\mathcal{P}(\mathcal{O}^{\dagger}_{\pm\Delta})^n|C^k_{N-k}\rangle$ which have all quantum numbers properly defined. In fact, the use of the $\mathcal{P}$ operator opens another possibility for $\mathcal{O}^{\dagger}$-type operators:

$$\Sigma^{\dagger}_{1} = \sum_{m} \sqrt{m+1}\, c^{\dagger}_{m+1,\downarrow} c_{m,\uparrow}. \tag{5}$$

acting on any state $|C^k_{N-k}\rangle$ does not have well defined M_{CM}, but $\mathcal{P}(\Sigma^{\dagger}_1)^n|C^k_{N-k}\rangle$ has $M_{CM} = 0$ for all n.

In order to check whether these ansatz states are eigenstates of Hamiltonian (1), we have performed numerical diagonalisations for $N \leq 9$. The overlap between $\mathcal{P}(\mathcal{O}^{\dagger}_{\pm\Delta})^n|C^k_{N-k}\rangle$ and the exact lowest energy state in the (M, S_z) subspace they belong to is always higher than 0.99. This is so for $\mathcal{O}$ being Υ, Λ, or Σ, and for all Δ, n, and k which satisfy the restrictions stated above on the applicability to these operators. There is only one case in which two different states of the form $\mathcal{P}(\mathcal{O}^{\dagger}_{\pm\Delta})^n|C^k_{N-k}\rangle$ belong to the same subspace: $\mathcal{P}(\Lambda^{\dagger}_1)^n|C^k_{N-k}\rangle$ and $\mathcal{P}(\Sigma^{\dagger}_1)^n|C^k_{N-k}\rangle$ We find that, for $k > 0$, the former one has always a high overlap with the exact lowest energy state of the subspace it belongs to, while the later one (when orthogonalised to the former one) has an overlap again larger than 0.99 with one of the low-lying excited states in that subspace. For $k = 0$ (i.e. $\nu = 1$), there are not states generated by $\Lambda^{\dagger}_1$, and $\mathcal{P}(\Sigma^{\dagger}_1)^n|C^k_{N-k}\rangle$ becomes the lowest energy state of the subspace with $M = N(N-1)/2 + n$ and $S_z = N/2 - n$.

Now we investigate the physical nature of the different kinds of states. Let us focus first on states of the form $\mathcal{P}(\Lambda^{\dagger}_1)^n|C^1_{N-1}\rangle$. Simpler wavefunctions can be obtained by the inverse transformation to the one which projects a skyrmionic BCS-like single Slater determinant into a state with well defined M. [7,10] We obtain

$$|\Psi(\varphi,\xi)\rangle = \sum_{n=0}^{N-2} \frac{e^{in\varphi}}{n!} \xi^n (\Lambda^{\dagger}_1)^n |C^1_{N-1}\rangle \tag{6}$$

$$= -c^{\dagger}_{0\downarrow} \prod_{j=0}^{N-2} \left(c^{\dagger}_{j\uparrow} + \frac{\xi}{\sqrt{j+1}} e^{i\varphi} c^{\dagger}_{j+1\downarrow}\right)|\phi\rangle \tag{7}$$

$|\phi\rangle$ being the vacuum state. From this form, $|\Psi(\varphi,\xi)\rangle$ is readily recognised as a HF wavefunction describing skyrmions with topological charge 1; [6,7,9] it depends on two parameters: φ is a variable fixing a broken-symmetry direction of the spin in the xy plane, while parameter ξ controls the skyrmion size (i.e. its M and S_z). From eq. (7), $S^{SK}_z(r) \sim (r^2 - 2\xi^2)/(r^2 + 2\xi^2)$ at large distances, as for skyrmions in the CNLSM, which implies a size [6] $\lambda = \xi/\sqrt{2}$. Remarkably, such dependence of $S^{SK}_z(r)$ holds even in the case of skyrmion sizes comparable with the magnetic length $l_B = \sqrt{\hbar/m^*\Omega}$, regime well beyond the range of applicability of the classical model. A technical point, of relevance in the interpretation of these states, is that in eq. (6) the projector $\mathcal{P}$ is not included. Otherwise $|\Psi(\varphi,\xi)\rangle$ would not be a BCS-like single Slater determinant. Corrections introduced by $\mathcal{P}$ are proportional to S_z^{-1}, so for $n << N$ and $N >> 1$ they are negligible. In the small N limit, or for $n \sim N/2$, the skyrmion is "deformed" and its identification as such becomes fuzzier. For instance, we find (for N even) $\mathcal{P}(\Lambda^{\dagger}_1)^{(N/2)-1}|C^1_{N-1}\rangle = \mathcal{P}(\Upsilon^{\dagger}_{-1})^{N/2}|C^0_N\rangle$.

By a similar argument, we identify excitations created by $\Lambda^{\dagger}_{\Delta}$ as skyrmions with topological charge Δ. Recall that, although skyrmions with $\Delta > 1$ are never the lowest energy excitation branch in this model, they are eigenstates appearing from partially-ferromagnetic compact systems with, at least, Δ electrons with spin $\downarrow$.

A physical identification of states generated by $\Sigma^{\dagger}_1$ or $\Upsilon^{\dagger}_{-\Delta}$ is not so direct. From the form of the operators both of them are *edge* spin-textures. Unlike what occurs for skyrmions where S_z changes sign as a function of r, their $S_z(r)$ is always positive. They resemble spin-waves in that they appear as excitations of any compact (partially-ferromagnetic) system.

In an infinite system, electron-hole symmetry can be invoked to obtain other spin-textures. In particular antiskyrmions arise as $(\Lambda^{\dagger}_{\Delta})$-type excitations of states $|C^{N\to\infty}_k\rangle$ of holes. For the antiskyrmion with topological charge -1 and $S_z = n$, we obtain the wavefunction $(\sum_{m=1}^{\infty} m^{-1/2} c^{\dagger}_{m-1,\downarrow} c_{m,\uparrow})^n c_{0,\uparrow}|C^0_{N\to\infty}\rangle$. For a finite system, the hole state complementary to the electron state $c_{0,\uparrow}|C^0_{N+1}\rangle$ is not a compact state and it has $M_{CM} \neq 0$. In this case, antiskyrmions cannot be built up by operators of the form $\mathcal{O}^{\dagger}_{\pm\Delta}$.

Numerical diagonalisations for small N show that, on increasing B, the spin-flip transition from $|C^{k+1}_{N-(k+1)}\rangle$ to $|C^k_{N-k}\rangle$ occurs through a ladder of states of the form $\mathcal{P}(\Lambda^{\dagger}_1)^n|C^{k+1}_{N-(k+1)}\rangle$ and $\mathcal{P}(\Upsilon^{\dagger}_{-1})^n|C^k_{N-k}\rangle$. In this letter we concentrate on the transition from $|C^1_{N-1}\rangle$ to $|C^0_N\rangle$, which is sketched in Fig. 1. Energies of skyrmion-like states $|\Psi^{SK}_n\rangle = \mathcal{P}(\Lambda^{\dagger}_1)^n|C^1_{N-1}\rangle$ take the form [19]

$$\frac{\langle\Psi^{SK}_n|H|\Psi^{SK}_n\rangle}{\langle\Psi^{SK}_n|\Psi^{SK}_n\rangle} = E^{SP}(M,S_z) + E_0 + \alpha_{\Lambda} n + \beta_{\Lambda} \begin{pmatrix} n \\ 2 \end{pmatrix} \tag{8}$$

where E_0 is the interaction energy of the initial compact state, $|C^1_{N-1}\rangle$ in this case. Both α_Λ and β_Λ have analytical expressions in terms of $V_{m_1m_2m_3m_4}$ which can be calculated, by using Wick's theorem. We find $\alpha_\Lambda < 0$ and $\beta_\Lambda > 0$, so skyrmions in a QHD behave as confined bosons interacting via a two-body repulsive interaction. Figure 2 shows the dispersion relation of the skyrmion branch for $N = 30$. Energies for $\mathcal{P}(\Upsilon^\dagger_{-1})^n|C^0_N\rangle$ have the form of eq.(8), with different expressions for E_0, α_Υ and β_Υ. We find that β_Υ is negligible for all N. The transition from $|C^1_{N-1}\rangle$ to $|C^0_N\rangle$ presents two cases:

i) for $N < N_0$, where N_0 depends on the Zeeman energy but is $N_0 \lesssim 10$, the transition is through a ladder of states with decreasing S_z first, until it reaches its minimun possible value and increasing S_z afterwards, as depicted in Fig. 1.

ii) for $N > N_0$ the minimum of the skyrmion branch is not at the end of the branch ($n = N/2 - 1$), as shown in Fig. 2 for $N = 30$. The transition is through all the skyrmion states with n less than a certain n^{SK}, which depends on g, going directly to $|C^0_N\rangle$ afterwards without passing through any of the $\mathcal{P}(\Upsilon^\dagger_{-1})^n|C^0_N\rangle$ states. n^{SK} labels the spin of the largest skyrmion which becomes GS when varying B.

The spin of the GS is a measurable quantity which may be controlled varying the Zeeman energy by means of a tilted B. Figure 3 shows n^{SK} as a function of Zeeman energy, calculated from $\Psi^{SK}_n\rangle$ and the HF approximation (HF energies can be obtained using eq. (6)). The HF approach gives slightly smaller values for n^{SK}. This is so because, although the HF dispersion relation is very accurate (see inset of Fig. 3(a) and note the small energy scale) its minimum is shifted to smaller n.

Experimental information on the transition from $|C^1_{N-1}\rangle$ to $|C^0_N\rangle$ is already available: single electron capacitance [11] and transport [13] experiments in quantum dots have been interpreted [13,14] in terms of highly correlated GS's appearing in numerical diagonalizations performed for $N \leq 5$. Here, we have identified those states as skyrmions and edge-spin excitations and shown that skyrmions appear for all N.

To complete our discussion on skyrmions we analyse the thermodinamic limit. In the $g = 0$ limit, CNLSM predicts that skyrmions in an infinite system have smaller interaction energy than $|C^1_{N\to\infty}\rangle$ by an amount $\Delta E_{int} = -\sqrt{\pi/32}(e^2/4\pi\epsilon l_B)$. [5] Figure 4 shows n^{SK} at $g = 0$ and the interaction energy of the skyrmion of size n^{SK} as a function of N. n^{SK} grows slowly with N, while ΔE_{int} tends to the classical value.

From numerical diagonalisations we know that, for small N, [15,19] only two kind of excitations become the new GS of the system when increasing B: charge magnetoexcitons $J^\dagger_\Delta|C^0_N\rangle$ [17] or edge spin excitations $\mathcal{P}(\Sigma^\dagger_1)^n|C^0_N\rangle$. However, for an infinite 2D system, antiskyrmions must be the new GS of the system for $\nu \lesssim 1$. Since we do not have analytical expresions for wavefunctions and energies of antiskyrmions in the intermediate N regime, we cannot conclude whether they are the new GS in this case. With this caveat in mind, we concentrate on charge and edge-spin excitations at $\nu = 1$. For arbitrary N, expressions for magnetoexciton energies have been reported. [17,21] We find that edge spin excitation energies are of the form (8) with negligible β_Σ. Therefore, their energies are linear in n, i.e. in M, see Fig. 2 (notice that this is equivalent to a dispersion relation which is quadratic in linear momentum k, as occurs for excitations produced by a broken continuous symmetry). Which type of excitations has lower energy depends on both N and Zeeman energy. In the $g = 0$ limit, we find that the edge spin branch has lower energy than the charge excitation branch for $N < N_{min}$ and vice versa for $N > N_{min}$ where $N_{min} \simeq 100$ for the Coulomb repulsion. On increasing Zeeman energy, the charge branch is favoured. For instance, for GaAs ($g = 0.44$), $N_{min} \sim 20$ for $\hbar\omega_0 = 2meV$ while for $\hbar\omega_0 = 5.4meV$ the charge branch has lower energy than the spin branch for any number of electrons ($N_{min} = 0$).

We thank L. Brey and A. H. MacDonald for useful discussions. Work supported by CICYT of Spain under contracts No. MAT 94-0982-C02 and 94-0058-C02.

---

[1] R. Tycko *et al.*, Science **268**, 1460 (1995); S. E. Barret *et al.*, Phys. Rev. Lett.,**74**, 5112 (1995).
[2] A. Schmeller *et al.*, preprint cond-mat/9506133.
[3] E. H. Aifer *et al.*, preprint cond-mat/9507069.
[4] E. H. Rezayi, Phys. Rev. B **36**, 5454 (1987).
[5] S. L. Sondhi *et al.*, Phys. Rev. B **47**, 16419 (1993).
[6] H. A. Fertig *et al.*, Phys. Rev. B **50**, 11018 (1994).
[7] K. Moon *et al.*, Phys. Rev. B **51**, 5138 (1995).
[8] X. C. Xie and Song He, Phys. Rev. B **51**, 5138 (1995).
[9] L. Brey *et al.*, Phys. Rev. Lett., **75**, 2562 (1995).
[10] C. Nayak and F. Wilczek, preprint cond-mat/9505081.
[11] R. C. Ashoori *et al.*, Phys. Rev. Lett. **71**, 613 (1993).
[12] O. Klein *et al.*, Phys. Rev. Lett. **74**, 785 (1995).
[13] T. Schmidt *et al.*, Phys. Rev. B **51**, 5570 (1995).
[14] J. J. Palacios *et al.*, Phys. Rev. B **50**, 5760 (1994).
[15] S.-R. Eric Yang *et al.*, Phys. Rev. Lett. **71**, 3194 (1993).
[16] R. K. Kamilla and J. K. Jain, Phys. Rev. B **52**, 2798 (1995).
[17] J. H. Oaknin *et al.*, Phys. Rev. Lett., **74**, 5120 (1995).
[18] S. A. Trugman and S. Kivelson, Phys. Rev. B **31**, 5280 (1985).
[19] Details of the calculation will be published elsewhere.
[20] $\mathcal{P}$ is generated from spin-ladder operators S^+ and S^-.
[21] L. Martin-Moreno *et al.*, Solid St. Elect., to be published.

FIG. 1. Sketch of M and S_z of the different states which, in some conditions, become GS of the QHD around $\nu = 1$. Compact (●), skyrmions (○), edge spin excitations (×), (⋄) and charge excitations (△) are shown.

FIG. 2. Dispersion relations of skyrmions (Λ_1), edge spin excitations (Υ_{-1}), (Σ) and charge excitations (J) for $N = 30$. ΔM is the excess angular momentum with respect $|C^1_{N-1}\rangle$. Parameters for $E^{SP}(M, S_z)$ are taken such that $|C^0_N\rangle(*)$ and $|C^1_{N-1}\rangle(\circ)$ have the same total energy.

FIG. 3. Exact (continous line) and HF (dotted line) S_z for the largest skyrmion which becomes GS when varying B (in the text refered to as n^{SK}) as a function of g for $N = 30$ and (a) $\hbar\omega_0 = 2meV$, (b)$\hbar\omega_0 = 5.4meV$. Inset in panel (a) shows the comparison between exact (dots) and HF (dotted line) total energy for skyrmions (in units of $e^2/(4\pi\epsilon l_B)$). Parameters for $E^{SP}(M, S_z)$ are taken as in figure 2.

FIG. 4. n^{SK} (dotted line) and interaction energy of the skyrmion of $S_z = n^{SK}$ with respect to that of $|C^1_{N-1}\rangle$ (continous line) as a function of N, in the $g = 0$ limit. $*$ marks the interaction energy for skyrmions given by the CNLSM in an infinite system. All energies are in units of $e^2/(4\pi\epsilon l_B)$.


S_z
$N/2$
$N/2-1$
$N/2-2$
$N/2-3$
$\vdots$
0
$|C^1_{N-1}\rangle$
$P\Lambda_1^+|C^1_{N-1}\rangle$
$P(\Lambda_1^+)^2|C^1_{N-1}\rangle$
$P(\Lambda_1^+)^{(N/2)-1}|C^1_{N-1}\rangle = P(\Upsilon_{-1}^+)^{N/2}|C^0_N\rangle$
$P(\Upsilon_{-1}^+)^2|C^0_N\rangle$
$P\Upsilon_{-1}^+|C^0_N\rangle$
$|C^0_N\rangle$
$J_1^+|C^0_N\rangle$
$J_2^+|C^0_N\rangle$
$P\Sigma_1^+|C^0_N\rangle$
$P(\Sigma_1^+)^2|C^0_N\rangle$
0
1
2
$\cdots$
$N/2-1$
$\cdots$
$N-3$
$N-2$
$N-1$
N
$N+1$
ΔM


N=30
Total energy
0.4
0.3
0.2
0.1
0.0
J
Σ
ϒ₋₁
Λ₁
0
5
10
15
20
25
30
35
40
45
Δ M


(a)
(b)
S_z
0
1
2
3
Total Energy
0.02
0.0
−0.02
ΔM
0
1
2
3
4
5
0.0
0.2
0.4
0.6
0.8
1.0
Lande g−factor



$n^{SK}\ (g=O)$
$-\Delta E_{int}$
N
0
200
400
600
800
1000
∞
2
4
6
8
10
12
14
0.3
0.32
0.34
0.36
0.38
0.4
0.42
0.44